

Recommended Childhood and Adolescent MSDH Immunization Schedule 2007†

Vaccine	Age ▶	Range of recommended ages			Catch-up Vaccination			Preadolescent assessment						
		Birth	1 mo	2 mos	4 mos	6 mos	12 mos	15 mos	18 mos	24 mos	4-6 years	11-12 years	13-14 years	15 years
Hepatitis B	HepB											HepB Series		
Inactivated Polio										IPV ⁴				
Diphtheria, Tetanus, Pertussis										DTaP	Tdap*	Tdap*		
Haemophilus Influenzae Type b														
Measles, Mumps, Rubella												MMR		
Varicella												Varicella		
Rotavirus#			RTV	RTV	RTV									
Meningococcal												MCV4	MCV4	
Pneumococcal			PCV	PCV	PCV							PCV	PPV	
Hepatitis A												HepA Series *		
Influenza												Influenza (yearly)		
Human Papillomavirus												HPV 3doses	HPV (3 doses)	

Pediarix™*
#1 #2 #3

PedvaxHIB™*
#1 #2

TRIHIBIT™*

ProQuad®*

Vaccines within broken line are for selected populations

†The complete ACIP/AAP/AAFP immunization schedule with appropriate footnotes can be accessed at <http://www.cdc.gov/mmwr/pdf/wk/mm5551-Immunization.pdf>

*Brand names are used for simplification and clarification

MDH clinics will use BOOSTRIX® for adolescent Tdap and VAQTA® for Hepatitis A for infants/young children

The minimum age for the first dose is 6 weeks. The first dose should be administered between 6 and 12 weeks of age (that is, until age 13 weeks). The 3rd dose should be administered through 32 weeks, (if not administered by this time, the third dose should be omitted).